CoLYTELY®/ GoLYTELY®/ NuLYTELY®/ TriLyte® (PEG) Instructions

Your doctor has prescribed PEG Bowel Prep Kit for your upcoming procedure. You may be feeling a little anxious, but don't worry, this instruction sheet will guide you through every step.

Access video instructions on our website.



Medication Guidelines

GLP-1 and phentermine medications

- Seven (7) days prior to your procedure, discontinue oral and injectable GLP-1 agonist, phentermine, and any weight loss medications, such as Bydureon BCise[®] (exenatide), Byetta[®] (exenatide), Mounjaro[®] (tirzepatide), Ozempic[®] (semaglutide), Rybelsus[®] (semaglutide), Saxenda[®] (liraglutide), Trulicity[®] (dulaglutide), Victoza[®] (liraglutide), Wegovy[®] (semaglutide), Zepbound (tirzepatide), Adipex, Fastin[®], Phentercot, Pro-Fast, Qsymia[®].
- Discontinue fiber and iron supplements, multivitamins with iron and high doses of fish oil.

Blood Thinning Medications

- Your GI provider will advise you whether to stop taking any blood thinning medications (anticoagulants and antiplatelets), after consulting with your prescribing physician.
- If you are currently taking aspirin, you may continue your dosage.

Diabetic Medications

 Three (3) days prior, diabetic patients, do not take SGLT2 medications, such as JARDIANCE[®] (empagliflozin), INVOKANA[®] (canagliflozin), FARXIGA[®] (dapagliflozin), SYNJARDY[®] (empagliflozin/metformin), or STEGLUJAN[®] (ertugliflozin/sitagliptin), etc.

Three Days Before Your Procedure

Clear Liquid Diet Guidelines

- For a successful bowel cleansing prep, it is essential that you **purchase** clear liquid items as indicated.
- Purchase your clear liquid diet items. For comfort, purchase diaper rash ointment, your preferred two-or-three-plied toilet paper, and/or flushable wet wipes.

Allowed Items

- ✓ Water
- Clear broths
- Coffee / tea without milk or creamers
- Clear juices without pulp
- ✓ Gatorade[®]
- Jell-0[®]
- Pulp-free Popsicles[®]

Items to Avoid

- × Dairy
- × Red or purple juices
- × Anything with pulp
- × Alcohol
- × Solid food

• Stop eating raw vegetables or vegetables containing seeds, corn, popcorn, nuts, and seeds.

Two Days Before Your Procedure

• Continue a low fiber diet and remember to avoid foods with nuts and seeds.

For Patients with Constipation

Please follow the magnesium citrate instructions. In the morning, begin a clear liquid diet and avoid solid foods until after your procedure. At 6:00 p.m., drink 10-ounces of magnesium citrate. You may mix with Gatorade or clear soda for flavor. If you have congestive heart failure or have kidney failure on dialysis (peritoneal or hemodialysis), do not drink and contact your gastroenterologist.

One Day Before Your Procedure

• Unless otherwise directed, begin your clear liquid diet in the morning and refrain from consuming solid food until after your procedure.



- In the morning, prepare the prep mixture by adding water to the indicated line (4 liters) near the top of the gallon jug. Shake until the powder is completely mixed with the water and chill in the refrigerator. The mixture cannot be mixed with any other liquid and must be consumed within 24 hours after mixing.
- At 6:00 p.m., start consuming the prep mixture by drinking an 8-ounce glass every 15-20 minutes until you have finished half of the prep mixture.
- Medication instructions for the evening before
 - Insulin dependent patients may take insulin if necessary, but at **half** the usual dose with close monitoring of glucose. Keep a sugary clear liquid available if necessary.
 - Non-insulin dependent diabetic diabetic patients, take your oral diabetic medication dose as usual.

 To comply with anesthesia requirements, do not use any tobacco products or marijuana on the day of your procedure. This includes smokeless tobacco, chewing tobacco, vapes, e-cigarettes, ZYN pouches, etc.

6 hours before the procedure



- Start the second dose of prep 6 hours before the procedure start time, following the same instructions as your first dose.
- You must finish the entire prep even if your bowels are already clear.
- You must finish your prep 4 hours before your procedure.

Medication instructions for the day of procedure

- Diabetic patients, follow these medication instructions.
 - Insulin dependent patients may take insulin if necessary, but at half the usual dose with close monitoring of glucose. Keep a
 sugary clear liquid available if necessary to regulate your blood sugar. Nothing by mouth 2 hours prior to your procedure.
 - Non-insulin dependent diabetic patients, hold your oral diabetic medication until after your procedure.
- Unless your doctor gave you other instructions, prescribed medication(s) may be taken as scheduled with a small sip of water at least <u>2 hours before your procedure</u>. Do not take anything else by mouth until your procedure is completed.
- Asthma patients, bring your inhaler to your appointment.
- It is imperative that you take all seizure and blood pressure medication on the morning of the procedure.

4 hours before the procedure

- Nothing by mouth until after your procedure.
- Do not chew gum or eat hard candy.
- You may brush your teeth but do not swallow.

Additional Day of Instructions

Because your procedure requires sedation, a driver (age 18 or over) must accompany you to your appointment. Your driver must sign in with you when you check in for your procedure and remain at the facility during the procedure to be available in case of an emergency; otherwise, you will not be able to proceed. You may not go home in a taxi, rideshare (Uber/Lyft), or take public transportation. The sedation you will receive will prevent you from driving safely or working the day of your procedure. Following your procedure, do not drive, operate heavy machinery, or sign legal documents for 24 hours.



Follow any additional instructions from your doctor. If you have questions, you can contact Gifthealth.

For additional information, scan the QR code for FAQs.

Need further assistance?

- Chat now on www.gifthealth.com
- **&** 833.614.2051
- care@gifthealth.com